

ARTICLE

RETRACTED ARTICLE: The decline of glomerular function is not always associated with the development of micro- and macroalbuminuria in hypertensive patients with type 2 diabetes

**R. Nosadini • A. Carboni • A. Manconi • F. Angius •
S. Caria • S. Cherchi • A. Satta • R. Faedda • D. Obinu •
M. Nieddu • A. Carraro • G. C. Tonolo**

Received: 7 January 2008 / Accepted: 21 May 2008
© The Author(s) 2008

This article has been withdrawn due to copyright infringement via plagiarism.

This article has been withdrawn due to copyright infringement via plagiarism.

R. Nosadini (✉)
Department of Endocrinology and Metabolic Diseases,
Internal Medicine Department, University of Sassari,
Viale San Pietro 8,
07100 Sassari, Italy
e-mail: noscia@tin.it

A. Carboni • A. Manconi • F. Angius • S. Caria • S. Cherchi •
D. Obinu • M. Nieddu • G. C. Tonolo (✉)
General Hospital ASL 2, Via Canova,
Olbia, Italy
e-mail: noscia@tin.it

A. Carraro
General Hospital Mirano,
Venice, Italy

A. Satta
Internal Medicine Department, University of Sassari,
Sassari, Italy

R. Faedda
Department of Nephrology, University of Sassari,
Sassari, Italy

Published online: 08 July 2008